

HOMOLOGICAL PROPERTIES OF COLOR LIE SUPERALGEBRAS

KENNETH L. PRICE

ABSTRACT. Let $\mathcal{L} = \mathcal{L}_+ \oplus \mathcal{L}_-$ be a finite dimensional color Lie superalgebra over a field of characteristic 0 with universal enveloping algebra $U(\mathcal{L})$. We show that $\text{gldim}(U(\mathcal{L}_+)) = \text{lFPD}(U(\mathcal{L})) = \text{rFPD}(U(\mathcal{L})) = \text{injdim}_{U(\mathcal{L})}(U(\mathcal{L})) = \dim(\mathcal{L}_+)$. We also prove that $U(\mathcal{L})$ is Auslander-Gorenstein and Cohen-Macaulay and thus that it has a QF classical quotient ring.

1. INTRODUCTION

Color Lie superalgebras are graded over an Abelian group G and generalize Lie superalgebras. Background information on Lie superalgebras and color Lie superalgebras can be found in [7] and [13] and [1] and [14], respectively. We will show, in a forthcoming paper [12], that the ideal structure of the enveloping algebra of a color Lie superalgebra can be very different from enveloping algebras of Lie superalgebras. By contrast, the results of this note illustrate that certain homological properties are the same. Our main strategy is to pass to the case where the grading group G is finitely generated, so that the color Lie superalgebra is determined by a Lie superalgebra and a 2-cocycle defined on G .

In §2 we define color Lie superalgebras and state a theorem due to Scheunert (see [14]). In §3 we calculate the finitistic and injective dimensions of the enveloping algebra of a finite dimensional color Lie superalgebra over a field of characteristic 0. The enveloping algebra of a finite dimensional color Lie superalgebra may have infinite global dimension (this is known for ordinary Lie superalgebras, see [2, Proposition 5]). However, for a finite dimensional color Lie superalgebra $\mathcal{L}$ which is *positively graded*, i.e. $\mathcal{L} = \mathcal{L}_+$, we prove that, in analogy with the nongraded case, $\text{gldim } U(\mathcal{L}) = \dim \mathcal{L}$. In particular, theorem 3.1 generalizes [8, Proposition 2.3].

In theorem 3.2 we show that the enveloping algebra $U(\mathcal{L})$ of a color Lie superalgebra is Auslander-Gorenstein and Cohen-Macaulay. It thus follows from [9, Theorem 1.4] that $U(\mathcal{L})$ has a (right and left) quasi-Frobenius (QF) classical quotient ring.

2. COLOR LIE SUPERALGEBRAS

Throughout k denotes a field of characteristic $\neq 2$, G denotes an Abelian group, and all algebras are associative k -algebras with 1. We call a map $\gamma : G \times G \rightarrow k^\times$

$(k^\times = k \setminus \{0\})$ a *skew-symmetric bicharacter* on G if it satisfies

- (1) $\gamma(f, gh) = \gamma(f, g)\gamma(f, h)$ and $\gamma(gh, f) = \gamma(g, f)\gamma(h, f)$ for any $f, g, h \in G$.
- (2) $\gamma(g, h)\gamma(h, g) = 1$ for any $g, h \in G$.

Note that (2) implies that $\gamma(g, g) = \pm 1$ for any $g \in G$. Set $G_\pm = \{g \in G : \gamma(g, g) = \pm 1\}$, then G_+ is a subgroup of G and $[G : G_+] \leq 2$. In [14, Lemma 2] it is shown that if G is finitely generated then the skew-symmetric bicharacters on G are completely determined by the 2-cocycles on G with coefficients in $k^\times$.

Generally, we shall be discussing objects which are graded over G so let $G\text{-vec}$ denote the category of G -graded vector spaces and graded linear maps. For any $V \in G\text{-vec}$, set $\partial x = g$ if $0 \neq x \in V_g$. Let γ be a skew-symmetric bicharacter on G . We will shorten our notation by writing $\gamma(x, y)$ instead of $\gamma(\partial x, \partial y)$ for homogeneous $0 \neq x \in V$, $0 \neq y \in W$ and $V, W \in G\text{-vec}$. Note that $G\text{-vec}$ is naturally contained in $\mathbb{Z}_2\text{-vec}$ via the decomposition $G = G_+ \dot{\cup} G_-$ and the group homomorphism $\pi : G \rightarrow \mathbb{Z}_2$ given by $\pi(G_+) = \bar{0}$ and $\pi(G_-) = \bar{1}$.

Definition. A (G, γ) -color Lie superalgebra is a Lie algebra a pair $(\mathcal{L}, \langle, \rangle)$ such that $\mathcal{L} \in G\text{-vec}$ and $\langle, \rangle : \mathcal{L} \otimes_k \mathcal{L} \rightarrow \mathcal{L}$ is a graded bilinear map which satisfies the following for any homogeneous $x, y, z \in \mathcal{L}$.

$$\begin{aligned} \gamma\text{-skew-symmetry} \quad & \langle x, y \rangle = -\gamma(x, y)\langle y, x \rangle \\ \gamma\text{-Jacobi identity} \quad & \gamma(z, x), \langle x, \langle y, z \rangle \rangle + \gamma(y, z), \langle z, \langle x, y \rangle \rangle + \gamma(x, y), \langle y, \langle z, x \rangle \rangle = 0 \end{aligned}$$

Example. (1) A Lie superalgebra is a $(\mathbb{Z}_2, \chi)$ -color Lie superalgebra where $\chi(\bar{i}, \bar{j}) = (-1)^{ij}$ for any $i, j \in \mathbb{Z}$.

- (2) If A is a G -graded algebra then there is a color Lie superalgebra structure A^- defined on A subject to the condition that $\langle a, b \rangle = ab - \gamma(a, b)ba$ for any nonzero homogeneous $a, b \in A$.

We shall now describe how to construct color Lie superalgebras from Lie superalgebras as in [14]. In as much as Lie superalgebras are $\mathbb{Z}_2$ -graded, we must consider G -gradings on Lie superalgebras which respect the $\mathbb{Z}_2$ -grading.

Definition. A Lie superalgebra $(L, [,])$, $L = L_{\bar{0}} \oplus L_{\bar{1}}$, is G -graded if $L = \bigoplus_{g \in G} L_g$, $[L_f, L_g] \subseteq L_{fg}$ for any $f, g \in G$, and $L_h \subseteq L_{\bar{0}}$ or $L_h \subseteq L_{\bar{1}}$ for each $h \in G$.

Starting with a Lie superalgebra L there are many choices of groups G and G -gradings on L . Let $G(L)$ be the smallest subgroup of G which grades L , that is, $G(L)$ is generated by $\{g \in G : L_g \neq 0\}$. We cannot always pass to the case that $G = G(L)$ since, for example, there are G -graded representations of L which are not $G(L)$ -graded.

Let $G(L)_+$ be the subgroup of $G(L)$ which is generated by $\{g \in G : U(L)_g \subseteq U(L)_{\bar{0}}\}$. Then $[G(L) : G(L)_+] \leq 2$. Set $G(L)_- = G(L) \setminus G(L)_+$; then the pair $(L, [,])$ is a $(G(L), \gamma_0)$ -color Lie superalgebra where γ_0 is the skew-symmetric bicharacter on $G(L)$ defined below.

$$\gamma_0(g, h) = \begin{cases} -1 & \text{if } g, h \in G(L)_- \\ 1 & \text{otherwise} \end{cases}$$

To continue our construction of a color Lie superalgebra from a Lie superalgebra, we need the notion of a 2-cocycle.

Definition. A 2-cocycle on G is a map $\sigma : G \times G \rightarrow k^\times$ which satisfies

$$\sigma(f, gh)\sigma(g, h) = \sigma(f, g)\sigma(fg, h)$$

for any $f, g, h \in G$.

If σ is a 2-cocycle on G then there is a skew-symmetric bicharacter γ defined on $G(L)$ by $\gamma(g, h) = \gamma_0(g, h)\sigma(g, h)\sigma(h, g)^{-1}$ for any $g, h \in G(L)$. Moreover, there is a $(G(L), \gamma)$ -color Lie superalgebra $(L^\sigma, [,]^\sigma)$ which has the same vector space structure as L but bracket $[,]^\sigma : L^\sigma \times L^\sigma \rightarrow L^\sigma$ defined subject to the condition that $[x, y]^\sigma = \sigma(\partial x, \partial y)[x, y]$ for any homogeneous $x, y \in L$.

By setting $f = h = e$ in Definition 2 (where e is the identity element of G) we obtain $\sigma(e, e) = \sigma(g, e) = \sigma(e, g)$ for any $g \in G$. Note that there is no loss in generality by assuming that $\sigma(e, e) = 1$ since we can replace σ with σ' , where $\sigma' = \sigma(e, e)^{-1}\sigma$ and have $L^\sigma \cong L^{\sigma'}$ as $(G(L), \gamma)$ -color Lie superalgebras.

Theorem 2.1 summarizes the results from [14] we are interested in. For the reader's convenience we will go over some basic definitions.

Definition. Let $(\mathcal{L}, \langle, \rangle)$ be a (G, γ) -color Lie superalgebra.

- (1) A linear map $\phi : \mathcal{L}_1 \rightarrow \mathcal{L}_2$ between (G, γ) -color Lie superalgebras $(\mathcal{L}_1, \langle, \rangle_1)$ and $(\mathcal{L}_2, \langle, \rangle_2)$ is called a *homomorphism* if $\phi(\langle x, y \rangle_1) = \langle \phi(x), \phi(y) \rangle_2$ for any $x, y \in \mathcal{L}_1$.
- (2) A G -graded representation of $\mathcal{L}$ is a pair (V, ρ) where $V \in G\text{-vec}$ and $\rho : \mathcal{L} \rightarrow \text{End}_k(V)^-$ is a homomorphism of color Lie superalgebras ($\text{End}_k(V)$ is a G -graded algebra since $V \in G\text{-vec}$).
- (3) For any G -graded algebra A , a graded map $\phi : \mathcal{L} \rightarrow A^-$ is called *compatible* if it is homomorphism of (G, γ) -color Lie superalgebras.
- (4) The *universal enveloping algebra* of $\mathcal{L}$ is a G -graded algebra $U(\mathcal{L})$ and a compatible map $\iota : \mathcal{L} \rightarrow U(\mathcal{L})$ which satisfies the property that for any compatible map $\phi : \mathcal{L} \rightarrow A$ there is a unique (graded) algebra homomorphism $\Phi : U(\mathcal{L}) \rightarrow A$ such that $\Phi \circ \iota = \phi$.

Theorem 2.1 (Scheunert). *Let $\mathcal{L}$ be a (G, γ) -color Lie superalgebra and L and L^σ as above.*

- (1) *If G is finitely generated, then any color Lie superalgebra can be obtained as an L^σ for appropriately chosen L and 2-cocycle σ .*
- (2) *The enveloping algebra $U(L^\sigma)$ is obtained from $U(L)$ by defining a new multiplication $*$ on U subject to the condition that for any homogeneous $x, y \in U(L)$ we have $x * y = \sigma(\partial x, \partial y)xy$.*
- (3) *For a graded representation $\rho : L \rightarrow \text{End}_k(V)$ of L , there is a graded representation $\rho^\sigma : L^\sigma \rightarrow \text{End}_k(V)$ of L^σ which is obtained from ρ subject to the condition that for any homogeneous $x \in L^\sigma$ and $v \in V$, $\rho^\sigma(x)(v) = \sigma(\partial x, \partial v)\rho(x)(v)$. This defines a category equivalence between the categories of graded representations of L and L^σ .*

We are particularly interested in parts (2) and (3) of theorem 2.1. More generally, consider an arbitrary G -graded algebra $\mathcal{R}$ and left $\mathcal{R}$ -modules V and W . An $\mathcal{R}$ -module homomorphism $\phi : V \rightarrow W$ is called *graded* if $\phi(V_h) \subseteq W_h$ for each $h \in G$. Let

${}_R\mathcal{M}^G$ denote the category whose objects are G -graded left $\mathcal{R}$ -modules and morphisms are all graded $\mathcal{R}$ -module homomorphisms from V to W , denoted $HOM_{\mathcal{R}}^0(V, W)$, where $V, W \in {}_R\mathcal{M}^G$. In particular, ${}_k\mathcal{M}^G = G\text{-}\mathbf{vec}$.

Lemma 2.2. *Let σ be a 2-cocycle on G and R a G -graded k -algebra.*

- (1) *There is a G -graded k -algebra R^σ with the same vector space structure as R and whose multiplication $*$ is obtained from the multiplication of R subject to the condition $r * s = \sigma(\partial r, \partial s)rs$ for any homogeneous $r, s \in R$.*
- (2) *For any $M \in {}_R\mathcal{M}^G$ there corresponds $M^\sigma \in {}_{R^\sigma}\mathcal{M}^G$ such that M^σ has the same vector space structure as M but the R^σ -module structure is obtained from the R -module structure of M subject to the condition $r.m = \sigma(\partial r, \partial m)rm$ for any homogeneous $r \in R$ and $m \in M$.*
- (3) *The functor $\sigma : {}_R\mathcal{M}^G \rightarrow {}_{R^\sigma}\mathcal{M}^G$ defined as in (2) is a category equivalence. In particular, $HOM_R^0(V, W) = HOM_{R^\sigma}^0(V^\sigma, W^\sigma)$. Moreover, if $V \in {}_R\mathcal{M}^G$ is graded free, then so is $V^\sigma \in {}_{R^\sigma}\mathcal{M}^G$.*
- (4) *For any $V, W \in {}_R\mathcal{M}^G$ and $\phi \in HOM_R^0(V, W)$ we have $\ker(\phi^\sigma) = (\ker \phi)^\sigma = \ker \phi$ and $\text{im}(\phi^\sigma) = (\text{im } \phi)^\sigma = \text{im } \phi$.*

Proof. It is easy to prove (1)-(3) directly. For (4), first note that $\ker \phi$, $\text{im } \phi$, $\ker(\phi^\sigma)$ and $\text{im}(\phi^\sigma)$ are G -graded submodules. For any homogeneous $x \in V$ we have $x \in \ker(\phi^\sigma) \Leftrightarrow x \in \ker(\phi) \Leftrightarrow x \in (\ker \phi)^\sigma$. This proves that $\ker(\phi^\sigma) = (\ker \phi)^\sigma = \ker \phi$. The proof that $\text{im}(\phi^\sigma) = (\text{im } \phi)^\sigma = \text{im } \phi$ is similar. ■

3. HOMOLOGICAL PROPERTIES OF $U(\mathcal{L})$

Throughout this section, γ is a skew-symmetric bicharacter on G and $G_+ = \{g \in G : \gamma(g, g) = \pm 1\}$. For any $V \in G\text{-}\mathbf{vec}$, set $V_\pm = \bigoplus_{g \in G_\pm} V_g$. If $V = V_+$ (respectively $V = V_-$) then V is called *positively* (respectively *negatively*) *graded*. We shall be primarily interested in the case where $\mathcal{L}$ is an Abelian color Lie superalgebra. If $\mathcal{L}$ is a positively graded, finite dimensional and Abelian (G, γ) -color Lie superalgebra, then $U(\mathcal{L}) \cong k^\gamma[\theta_1, \theta_2, \dots, \theta_n]$, the *color polynomial ring* in $n = \dim \mathcal{L}$ variables (see [1]). If $\mathcal{L}$ is a negatively graded, finite dimensional and Abelian (G, γ) -color Lie superalgebra, then $U(\mathcal{L}) \cong \Lambda^\gamma(\mathcal{L})$, the *color exterior algebra* (or *color Grassmann algebra*) of $\mathcal{L}$ (see [1]).

Example.

- (1) Choose $q \in k^\times$. Let γ be the skew-symmetric bicharacter on $G = \mathbb{Z}^2$ defined by $\gamma(g, h) = q^{g_1 h_2 - g_2 h_1}$ for any $g = (g_1, g_2)$, $h = (h_1, h_2) \in G$. Then $G = G_+$. Consider the Abelian (G, γ) -color Lie superalgebra with homogeneous basis $\{x, y\}$ such that $\partial x = (1, 0)$ and $\partial y = (0, 1)$. Then $\mathcal{L}$ is positively graded and $U(\mathcal{L}) \cong k[x, y : xy = qyx]$, the so-called *quantum plane*.
- (2) Choose $0 \neq q \in k^\times$. Let γ be the skew-symmetric bicharacter on $G = \mathbb{Z}^2$ defined by $\gamma(g, h) = (-1)^{g_1 h_1 + g_2 h_2} q^{g_1 h_2 - g_2 h_1}$ for any $g = (g_1, g_2)$, $h = (h_1, h_2) \in G$. Then $G_+ = \{(i, j) \in G : i + j \in 2\mathbb{Z}\}$ and $G_- = \{(i, j) \in G : i + j - 1 \in 2\mathbb{Z}\}$. Consider the Abelian (G, γ) -color Lie superalgebra with homogeneous basis $\{x, y\}$ such that $\partial x = (1, 0)$ and $\partial y = (0, 1)$. Then $\mathcal{L}$ is negatively graded and $U(\mathcal{L}) \cong k[x, y : xy = qyx]/I$, where I is the ideal generated by $\{x^2, y^2\}$.

If $V, W \in G\text{-vec}$ then the tensor product $V \otimes W \in G\text{-vec}$ has grading defined by $(V \otimes W)_g = \bigoplus_{h \in G} (V_h \otimes W_{gh^{-1}})$. For G -graded algebras A and B there is a γ -graded tensor product $A \widehat{\otimes}_k B$ with multiplication defined subject to the condition that $(a \widehat{\otimes} b)(a' \widehat{\otimes} b') = \gamma(b, a')aa' \widehat{\otimes} bb'$ for any nonzero homogeneous $a, a' \in A$ and $b, b' \in B$. The γ -graded tensor product is essential to the study of enveloping algebras of color Lie superalgebras. For example, the enveloping algebra of a color Lie superalgebra is not a Hopf algebra in the usual sense; it is only a Hopf algebra with respect to the γ -graded tensor product (see [1, §3.2.9] for details).

For any Noetherian ring R , the left (respectively right) finitistic dimension of R , denoted $\text{lFPD}(R)$ (respectively $\text{rFPD}(R)$), is the supremum of the projective dimensions of left (respectively right) R -modules of finite projective dimension. The following theorem generalizes [8, Proposition 2.3]. As in [8] and [9], we assume that $\text{char } k = 0$ in all that follows.

Theorem 3.1. *Let $\mathcal{L} = \mathcal{L}_+ \oplus \mathcal{L}_-$ be a finite dimensional color Lie superalgebra. Then $\text{gldim}(U(\mathcal{L}_+)) = \text{lFPD}(U(\mathcal{L})) = \text{rFPD}(U(\mathcal{L})) = \text{injdim}_{U(\mathcal{L})}(U(\mathcal{L})) = \text{dim}(\mathcal{L}_+)$.*

Proof. Let $U = U(\mathcal{L})$ denote the universal enveloping algebra of $\mathcal{L}$ and set $n = \text{dim}(\mathcal{L}_+)$. We first show that $\text{lFPD}(U) = \text{rFPD}(U) = \text{injdim}_U(U) = n \geq \text{gldim}(U(\mathcal{L}_+))$. Then we show that $\text{gldim}(U(\mathcal{L}_+)) \geq n$, that is, there exists a $U(\mathcal{L}_+)$ -module with projective dimension at least n . As in the case of ordinary Lie algebras, the existence of such a module will be demonstrated using the *Chevalley-Eilenberg complex* (see Chapter 7 of [16] for background).

1) With respect to the standard filtration, $gr(U)$ is Noetherian.

The filtration is defined by $U^{-1} = \{0\}$, $U^0 = k$ and, for $m > 0$, U^m is spanned by all monomials of length $\leq m$. By [1, Proposition III.2.8] U is Noetherian and the associated graded algebra $gr(U)$ is isomorphic to $k^\gamma[\theta_1, \theta_2, \dots, \theta_n] \widehat{\otimes} \Lambda^\gamma(\mathcal{L}_-)$, where $\Lambda^\gamma(\mathcal{L}_-)$ is the color exterior algebra determined by the vector space $\mathcal{L}_-$ with an Abelian color Lie superalgebra structure. Thus $gr(U)$ is an iterated Ore extension of $\Lambda = \Lambda^\gamma(\mathcal{L}_-)$ without derivations, that is, $gr(U) \cong \Lambda[\theta_1; \alpha_{\partial\theta_1}][\theta_2; \alpha_{\partial\theta_2}] \cdots [\theta_n; \alpha_{\partial\theta_n}]$. Therefore, $gr(U)$ is Noetherian by [11, Theorem I.2.10].

2) $\text{injdim}_U U \leq \text{injdim}_{gr(U)} gr(U) = n$ and $\text{gldim } U(\mathcal{L}_+) \leq \text{gldim } gr(U(\mathcal{L}_+)) = n$

Since $U^m = 0$ when $m < 0$, the filtration $\{U^m\}$ satisfies the *closure condition* of [3, §1.2.11]. Therefore, we may apply [3, Theorem 1.3.12] to obtain $\text{injdim}_U U \leq \text{injdim}_{gr(U)} gr(U)$. Moreover, $\text{gldim } U(\mathcal{L}_+) \leq \text{gldim } gr(U(\mathcal{L}_+))$ by [11, 7.6.18]. For any ring R and automorphism σ of R , one can show that $\text{gldim}(R[t; \sigma]) = \text{gldim}(R) + 1$ and $\text{injdim}_{R[t; \sigma]}(R[t; \sigma]) = \text{injdim}_R(R) + 1$ by similar methods to the proofs in the case that σ is the identity on R . Since $gr(U(\mathcal{L}_+)) \cong k^\gamma[\theta_1, \theta_2, \dots, \theta_n]$ we have $\text{gldim } gr(U(\mathcal{L}_+)) = n = \text{dim } \mathcal{L}_+$. By [6, Corollary 6.3] Λ is a Frobenius algebra; hence $\text{injdim}_{gr(U)} gr(U) = n = \text{dim } \mathcal{L}_+$ as $gr(U) \cong \Lambda^\gamma[\theta_1, \theta_2, \dots, \theta_n]$.

3) $\text{lFPD}(U) = \text{rFPD}(U) = \text{injdim}_U(U) \geq \text{gldim}(U(\mathcal{L}_+))$

Since $\text{gldim } U(\mathcal{L}_+)$ is finite and U is free over $U(\mathcal{L}_+)$ on both sides, we may apply [4, Theorem 1.4] to obtain $\text{gldim } U(\mathcal{L}_+) \leq \text{lFPD } U$. By [8, Proposition 2.1] we have $\text{injdim}_U U = \text{lFPD}(U) = \text{rFPD}(U)$.

The rest of the proof is dedicated to showing that if $\mathcal{L}$ is a positively graded finite-dimensional (G, γ) -color Lie superalgebra then $n = \dim \mathcal{L} \leq \text{gldim } U(\mathcal{L})$.

4) Pass to the case that $\mathcal{L} = L^\sigma$ for some Lie algebra L .

The modules we will consider are graded over the subgroup $G(\mathcal{L})$ of G generated by $\{g \in G : \mathcal{L}_g \neq 0\}$. Therefore, we can pass to the case that G is finitely generated by assuming that $G = G(\mathcal{L})$. By theorem 2.1(1), there is a G -graded Lie algebra L with bracket $[\cdot, \cdot]$ and a 2-cocycle σ on G such that $\mathcal{L} = L^\sigma$.

5) The Chevalley-Eilenberg complex $V_*(L) \xrightarrow{\varepsilon} k$ is a complex in $U(L)\mathcal{M}^G$.

We must show that $V_*(L) \xrightarrow{\varepsilon} k$ is a *graded complex*, i.e., the modules are graded and all differentiations are graded maps. Let ΛL denote the exterior algebra of L and $\Lambda^i L$ denote the i^{th} exterior algebra of L . Then $\Lambda L = \bigoplus_{i=1}^n \Lambda^i L$ can be G -graded so that the homogeneous component of degree $g \in G$ is $(\Lambda L)_g = \sum_{i=1}^n (\Lambda^i L)_g$ and $(\Lambda^i L)_g = \sum_{g_1 g_2 \dots g_i = g} L_{g_1} \wedge L_{g_2} \wedge \dots \wedge L_{g_i}$ where the sum is indexed by all appropriate $g_1, g_2, \dots, g_i \in G$.

We have $V_i(L) = U(L) \otimes_k \Lambda^i L$ which is a graded free left $U(L)$ -module such that $V_i(L)_g = \sum_{h \in G} (U(L)_h) \otimes_k (\Lambda^i L)_{hg^{-1}}$ so we only need to check that the *augmentation map* $\varepsilon : V_0(L) \rightarrow k$ is graded and, for $1 \leq i \leq n$, $d_i : V_i(L) \rightarrow V_{i-1}(L)$ is graded. The augmentation map $\varepsilon : V_0(L) = U(L) \rightarrow k$ is graded since it is induced from the Lie algebra map $L \rightarrow k$ which sends everything to zero, a graded map. It is also easy to see that $d : V_1(L) \rightarrow V_0(L)$ is graded since it is just the product map $u \otimes x \mapsto ux$. For $1 < i \leq n$, the map $d_i : V_i(L) \rightarrow V_{i-1}(L)$ is defined by the formula $d_i(u \otimes x_1 \wedge x_2 \wedge \dots \wedge x_i) = \theta_1 + \theta_2$ where θ_1 and θ_2 are defined below.

$$\begin{aligned} \theta_1 &= \sum_{j=1}^i (-1)^{j+1} u x_j \otimes x_1 \wedge x_2 \wedge \dots \wedge \hat{x}_j \wedge \dots \wedge x_i \\ \theta_2 &= \sum_{l < m} (-1)^{l+m} u \otimes [x_l, x_m] \wedge x_1 \wedge x_2 \wedge \dots \wedge \hat{x}_l \wedge \dots \wedge \hat{x}_m \wedge \dots \wedge x_i \end{aligned}$$

Therefore d_i is graded since if $u, x_1, x_2, \dots, x_i$ are homogeneous elements then $u \otimes x_1 \wedge x_2 \wedge \dots \wedge x_i, \theta_1$ and θ_2 are all homogeneous of the same degree.

6) The Chevalley-Eilenberg complex $V_*(\mathcal{L}) \xrightarrow{\varepsilon} k$ is a complex in $U\mathcal{M}^G$.

By parts (2), (3) and (4) of Lemma 2.2, there is a projective resolution $V_*(\mathcal{L})$ of $k = k^\sigma$ over $U(\mathcal{L})$ such that $V^i(\mathcal{L}) = V^i(L)^\sigma$. For $1 \leq i \leq n$ the module $V^i(L)$ is a graded free $U(L)$ -module hence the module $V^i(L)^\sigma$ is a graded free $U(L^\sigma)$ -module by Lemma 2.2(3).

7) $\text{gldim } U(\mathcal{L}) \geq n$

As in [16, Exercise 7.7.2], there is a graded representation $\rho : L \rightarrow \text{End}_k(\Lambda^n L)$ defined as below for $0 \neq y \in L$ and $x_1, x_2, \dots, x_n$ a (homogeneous) basis of L .

$$\rho(y)(x_1 \wedge x_2 \wedge \dots \wedge x_n) = \sum_{i=1}^n x_1 \wedge x_2 \wedge \dots \wedge [y, x_i] \wedge \dots \wedge x_n$$

Let $M = \Lambda^n L$ be the (graded) $U(L)$ -module defined by the above action. Then according to [16], $\text{Ext}_{U(L)}^n(k, M) \cong \ker(d_n^*)/\text{im}(d_{n-1}^*) \cong k$. By parts (2) and (4) of Lemma 2.2 and 6) above, we have $\ker d_n^* = (\ker d_n^*)^\sigma = \ker(d_n^*)^\sigma$ and $\text{im } d_{n-1}^* =$

$(\operatorname{im} d_{n-1}^*)^\sigma = \operatorname{im}(d_{n-1}^*)^\sigma$. This implies that

$$\operatorname{Ext}_U^n(k, M^\sigma) \cong \left(\operatorname{Ext}_{U(L)}^n(k, M) \right)^\sigma \cong k$$

therefore $\operatorname{pd}_U(M^\sigma) \geq n$ hence $\operatorname{gldim} U \geq n$ as desired.

The result follows from 3) and 7) above. ■

Theorem 3.2. *Let $\mathcal{L}$ be a finite dimensional color Lie superalgebra. Then $U(\mathcal{L})$ is Auslander-Gorenstein and Cohen-Macaulay and thus has a quasi-Frobenius classical quotient ring.*

Proof. We have Λ is Auslander-Gorenstein and Cohen-Macaulay since Λ is a Frobenius algebra (see [6, Corollary 6.3]). As $\operatorname{gr}(U(L))$ is an iterated Ore extension of Λ where each iteration is of the form $R[x; \sigma]$, it follows from [10, Lemma (ii), p. 184] and [5, Theorem 4.2] that $\operatorname{gr}(U(L))$ is Auslander-Gorenstein and Cohen-Macaulay. We thus conclude that $U(\mathcal{L})$ is Auslander-Gorenstein and Cohen-Macaulay by [3, Theorem 1.4.1] and [15, Lemma 4.4] (note that Theorem 3.1 is needed so that $\operatorname{injdim}_{U(\mathcal{L})} U(\mathcal{L}) < \infty$). The last statement follows from [9, Theorem 1.4]. ■

Acknowledgements

The author is indebted to Jim Kuzmanovich for many helpful conversations concerning this material.

REFERENCES

- [1] Yu. A. Bahturin, A. A. Mikhalev, V. M. Petrogradsky and M. V. Zaiciv, *Infinite Dimensional Lie Superalgebras*, DeGruyter, Berlin, 1992.
- [2] E. J. Behr, ‘Enveloping Algebras of Lie Superalgebras’, *Pacific J. Math.* **130** (1987), 9-25.
- [3] J-E Björk, ‘Filtered Noetherian Rings’, in *Noetherian Rings and Their Applications*, Math. Surveys and Monographs, vol. **24**, 58-97, Amer. Math. Soc., Providence, RI, 1987.
- [4] B. Cortzen, ‘Finitistic Dimension of Ring Extensions’, *Comm. Algebra* **10** (1982), 993-1001.
- [5] E. K. Ekström, ‘The Auslander Condition on Graded and Filtered Noetherian Rings’, in *Sem. Dubreil-Malliavin*, Lecture Notes in Math. **1404**, Springer-Verlag (1989), 220-245.
- [6] D. Fischman, S. Montgomery and H.-J. Schneider, ‘Frobenius Extensions of Subalgebras of Hopf Algebras’, to appear in *Trans. Amer. Math. Soc.*
- [7] V. G. Kac, ‘Lie Superalgebras’, *Adv. in Math.* **26** (1977), 8-96.
- [8] E. Kirkman, J. Kuzmanovich and L. Small, ‘Finitistic Dimensions of Noetherian Rings’, *Journal of Algebra* **147**(2) (1992), 350-364.
- [9] E. Kirkman and J. Kuzmanovich, ‘Minimal Prime Ideals in Enveloping Algebras of Lie Superalgebras’, *Proc. Amer. Math. Soc.* **124**(6), 1693-1702.
- [10] T. Levasseur and J. T. Stafford, ‘The Quantum Coordinate Ring of the Special Linear Group’, *J. Pure Appl. Alg.* **86** (1993), 181-186.
- [11] J. C. McConnell and J. C. Robson, *Noncommutative Noetherian Rings*, Wiley, Chichester, England, 1987.
- [12] K. L. Price, ‘Primeness Criteria for Universal Enveloping Algebras of Lie Color Algebras’, manuscript in preparation.
- [13] M. Scheunert, *The Theory of Lie Superalgebras: An Introduction*, Springer Lecture Notes in Mathematics (716), 1979.
- [14] M. Scheunert, ‘Generalized Lie Algebras’, *J. Math. Phys.* **20**(4) (1979), 712-720.
- [15] J. T. Stafford and J. J. Zhang, ‘Homological Properties of (Graded) Noetherian PI Rings’, *J. Algebra* **168** (1994), 988-1026.
- [16] C. A. Weibel, *An Introduction to Homological Algebra*, Cambridge Studies in Advanced Mathematics (38), Cambridge University Press, 1994.